

Tilburg University

Analyzing simulation experiments with common random number, part II

Kleijnen, J.P.C.

Publication date:
1988

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):

Kleijnen, J. P. C. (1988). *Analyzing simulation experiments with common random number, part II: Rao's approach*. (Research memorandum / Tilburg University, Department of Economics; Vol. FEW 332). Unknown Publisher.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

CBM
R

7626
1988
332


UNIVERSITY
HOLIERE
UNIVERSITEIT
BRABANT

POSTBOX 90153
5000 LE TILBURG
THE NETHERLANDS



DEPARTMENT OF ECONOMICS
RESEARCH MEMORANDUM

7626
1988
332



ANALYZING SIMULATION EXPERIMENTS WITH
COMMON RANDOM NUMBER, PART II:
RAO'S APPROACH

Jack P.C. Kleijnen

FEW 332

R46

518.931

June 1988

ANALYZING SIMULATION EXPERIMENTS WITH COMMON RANDOM NUMBERS, PART II:
RAO'S APPROACH

JACK P.C. KLEIJNEN

School of Business and Economics, Catholic University Brabant
(Katholieke Universiteit Brabant), 5000 LE Tilburg, Netherlands

Abstract

This note uses Rao (1959)'s approach based on Hotelling's statistic, to analyze a linear regression model with correlated responses. Correlated responses arise if a simulation model uses common random numbers. Rao's approach is compared to Kleijnen (1988)'s approach. Rao's lack of fit test is superior, which is quantified by a Monte Carlo study. Confidence intervals for individual regression parameters differ only slightly, when using Rao's and Kleijnen's approach respectively.

1. Introduction

Recently Kleijnen (1988) discussed the analysis of simulation experiments with common random number; Table 1 is reproduced from Kleijnen (1988, p. 66). These simulation data are analyzed through a linear regression model $y = X\beta + e$ with a nondiagonal covariance matrix. In this note we reconsider the analysis of this model, applying Rao (1959).

TABLE 1
Data of Simulation Experiment

Factor combination (effects) ($\beta_1 \dots \beta_Q$)	Replicated responses (seed 1) (seed 2) ... (seed m)				Average response	Estimated (co)variances
$x_{11} \dots x_{1Q}$	y_{11}	y_{12}	\dots	y_{1m}	\bar{y}_1	$\hat{\sigma}_1^2 \hat{\sigma}_{12} \dots \hat{\sigma}_{1n}$
$x_{21} \dots x_{2Q}$	y_{21}	y_{22}	\dots	y_{2m}	\bar{y}_2	$\hat{\sigma}_2^2 \dots \hat{\sigma}_{2n}$
$x_{i1} \dots x_{iQ}$	y_{i1}	y_{i2}	\dots	y_{im}	\bar{y}_i	$\hat{\sigma}_i^2 \dots \hat{\sigma}_{in}$
$x_{n1} \dots x_{nQ}$	y_{n1}	y_{n2}	\dots	y_{nm}	\bar{y}_n	$\hat{\sigma}_n^2$

Table 1 shows that there are m independent observations on the n -variate vector $\mathbf{y} = (y_1, \dots, y_i, \dots, y_n)'$ which yields the following unbiased estimators of $\sigma_{ii'} = \text{cov}(y_i, y_{i'})$:

$$\hat{\sigma}_{ii'} = \frac{\sum_{r=1}^m (y_{ir} - \bar{y}_i)(y_{i'r} - \bar{y}_{i'})}{m-1} \quad (i, i' = 1, \dots, n) \quad (m \geq 2), \quad (1.1)$$

where $\bar{y}_i = \sum_{r=1}^m y_{ir}/m$. We do not assume a specific pattern for the covariance matrix; all we assume is that the covariance matrix $\mathbf{Q}_y = (\sigma_{ii'})$ is non-singular. Kleijnen (1988, p. 67) proposed two different point estimators for the Q effects $\beta = (\beta_j)$. The Ordinary Least Squares (OLS) estimator is

$$\hat{\beta} = (\mathbf{X}'\mathbf{X})^{-1}\mathbf{X}'\bar{\mathbf{y}} \quad (1.2)$$

where $\mathbf{X} = (x_{ij})$ with $j = 1, \dots, Q$, and $\bar{\mathbf{y}} = (\bar{y}_i)$; also see Table 1. The Estimated Generalized Least Squares (EGLS) estimator is

$$\hat{\tilde{\beta}} = (\mathbf{X}'\hat{\mathbf{Q}}_y^{-1}\mathbf{X})^{-1}\mathbf{X}'\hat{\mathbf{Q}}_y^{-1}\bar{\mathbf{y}}, \quad (1.3)$$

where $\hat{\Omega}_y = (\hat{\sigma}_{ii})$ is assumed to be non-singular. The corresponding estimated covariance matrices of the estimators for β are

$$\hat{\Omega}_{\hat{\beta}} = (X'X)^{-1}X'\hat{\Omega}_yX(X'X)^{-1}/m \quad (1.4)$$

and

$$\hat{\Omega}_{\hat{\beta}} \approx (X'\hat{\Omega}_y^{-1}X)^{-1}/m \quad (1.5)$$

where \approx means that (1.5) holds only asymptotically ($m \geq 25$?; see Kleijnen, 1988, p. 68).

2. Kleijnen (1988)'s versus Rao (1959)'s validation test

Thirty years ago Rao considered a related problem, namely multivariate linear regression analysis. Unfortunately he did not use the symbols that have become standard in regression analysis. In the following paragraph we shall first give his formulas and then translate them into the symbols of Kleijnen (1988). (Readers interested in the resulting formula, can skip to equation 2.4.b).

Rao (1959, p. 49) considers the linear model

$$E(y) = A\tau \quad (2.1.a)$$

TABLE 2

Symbols in Rao (1959) and Kleijnen (1988) in order of appearance

Rao	:	A	τ	p	m	\wedge	y	e	S	f	r	$\hat{\tau}$	i
Kleijnen:	X	β	n	Q	Ω_y	\bar{y}	m	$\hat{\Omega}_y$	m-1	n-Q	$\hat{\beta}$	j	

where $y = (y_1, \dots, y_p)'$, A is the $p \times m$ matrix of coefficients, and τ is the vector of m regression parameters. Kleijnen (1988, p. 67) uses the model

$$E(y) = E(\bar{y}) = X\beta \quad (2.1.b)$$

Let $x := y$ mean that Rao's symbol x becomes Kleijnen's symbol y . So $A := X$, $\tau := \beta$, $p := n$, $m := Q$; also see Tables 1 and 2. Rao (1959, p. 49) further assumes "The dispersion matrix of y_1, \dots, y_p is $e^{-1}\Lambda$ where e is a known constant and Λ is unknown, but an estimator S of Λ is available and has Wishart's distribution based on f degrees of freedom independently of y_1, \dots, y_p . The matrix Λ is assumed to be non-singular". Analogously, Kleijnen (1988, p. 67) assumes that the covariance matrix of y_1, \dots, y_n is Ω_y and the covariance matrix of the averages $\bar{y}_1, \dots, \bar{y}_n$ is $\Omega_{\bar{y}} = \frac{1}{m} \Omega_y$. So if $\Lambda := \Omega_y$ and $y := \bar{y}$ then $e := m$. To check this translation we note that Rao's equation (2.6)

$$\min e(y-A\tau)' \Lambda^{-1} (y-A\tau) \quad (2.2.a)$$

corresponds to Kleijnen's EGLS criterion

$$\min (\bar{y}-X\beta)' \Omega_{\bar{y}}^{-1} (\bar{y}-X\beta) . \quad (2.2.b)$$

We note that Rao's example leading to his equation (1.2) is wrong: $e = 1/n$ should be $e = n$. We further have $S := \hat{\Omega}_y$ with degrees of freedom $f := m-1$. Next Rao (1959, p. 50) introduces " r equal to p minus the rank of A ". So $r = p - \text{rank}(A) := n - \text{rank}(X)$; usually $r := n-Q$; for saturated designs we have, by definition, $r := 0$. To test the model specification Rao uses Hotelling's statistic T_r which according to his equations (2.4) and (2.9) equals

$$T_r = \frac{e}{f} \min(y-A\tau)' S^{-1} (y-A\tau) . \quad (2.3.a)$$

Using (2.3) we get in Kleijnen's symbols:

$$T_r = \frac{1}{(m-1)} (\bar{y} - \hat{X}\hat{\beta})' \hat{Q}_y^{-1} (\bar{y} - \hat{X}\hat{\beta}) \quad (2.3.b)$$

Rao's equation (2.5) states that Hotelling's statistic T_r corresponds to the F statistic:

$$F_{r, f+1-r} = \frac{f+1-r}{r} T_r \quad (2.4.a)$$

or in Kleijnen's symbols, assuming for simplicity of notation that $\text{rank}(X) = Q$,

$$F_{n-Q, m-n+Q} = \frac{m-n+Q}{n-Q} \frac{1}{(m-1)} (\bar{y} - \hat{X}\hat{\beta})' \hat{Q}_y^{-1} (\bar{y} - \hat{X}\hat{\beta}) \quad (2.4.b)$$

We can interpret this equation as follows. A perfect fit ($\bar{y} = \hat{X}\hat{\beta}$) means that $T_r = 0$ and hence $F = 0$ and we do not reject the specified model; high estimated variances \hat{Q}_y^2 mean that large residuals ($\bar{y} - \hat{X}\hat{\beta}$) are accepted. Note that, if m goes to infinity then $F_{n-Q, m-n+Q}$ approaches χ_{n-Q}^2 . The F test exists only if $n > Q$ (non-saturated design).

The F statistic of (2.5.b) is a generalization of the F test for lack of fit in the classical experimental design literature, which assumes $\hat{Q}_y = \sigma^2 I$. The F test compares the estimated residuals (lack of fit) to the pure estimated noise \hat{Q}_y . Kleijnen (1988, p. 71) proposes an alternative approach: estimate β from one set of simulation data; use this estimate to predict the simulation output for a new input combination; compare the predictor \hat{y}_{n+1} to the observed simulation output y_{n+1} , using a t statistic. Kleijnen (1988, p. 71) states: "This test is simplest if we make \hat{y}_{n+1} ... and y_{n+1} independent, i.e., if we use a new seed for y_{n+1} ". However, such an approach excludes cross-validation: if $n > Q$ then cross-validation means that we delete one factor combination i ($i=1, \dots, n$) and estimate β from the remaining $n-1$ combinations ($X_{-i}, \bar{y}_{-i}, \hat{Q}_{y(-i)}$); the estimate for β is used to compute the OLS predictor $\hat{y}_i(\hat{\beta}_{-i})$ and/or the EGLS predictor $\hat{y}_i(\hat{\beta}_{-i})$; this predictor is compared to the simulation response \bar{y}_i , using Studentization, i.e., dividing the prediction error $\hat{y}_i - \bar{y}_i$ by its standard deviation $(\text{var}(\hat{y}_i - \bar{y}_i))^{\frac{1}{2}}$. If no common random numbers are used, this standard deviation is simple to derive; see Kleijnen (1983). If, however,

common random numbers are used, then the simulation responses y_i and y_i' are correlated, and the question arises how sensitive this t test is to this correlation. Before we try to answer that question, we consider another aspect of the regression analysis.

3. Kleijnen versus Rao's confidence intervals for β

Both Rao (1959, pp. 56-57) and Kleijnen (1987, p. 218) state that tests for individual parameters β_j are relevant, only after the specification of the regression model is tested (see § 2). To obtain confidence intervals for the parameters, Rao proceeds as follows (actually he considers the more general problem of a set of linear hypotheses).

Rao (1959, p. 52) introduces $C = (A'S^{-1}A)^{-1} := (X'\hat{Q}_yX)^{-1}$ which equals $m\hat{\hat{\beta}}$ asymptotically; see (1.5). Rao (1959, p. 53) defines c_{ii} as the

i -th diagonal element of C ; hence $c_{ii} := m \hat{\text{var}}(\hat{\beta}_j)$ asymptotically. His eq. (4.4) gives the following confidence limits to τ_i :

$$\hat{\tau}_i \pm t \left[\frac{e(f-r)}{c_{ii} f(1+T_r)} \right]^{-\frac{1}{2}} \quad (3.1.a)$$

where t has $f-r$ degrees of freedom; in our symbols this becomes

$$\hat{\beta}_j \pm t_v \left[\frac{(m-1-n+Q)}{\hat{\text{var}}(\hat{\beta}_j)(m-1)(1+T_r)} \right]^{-\frac{1}{2}} \quad (3.1.b)$$

where $v = m-1-n+Q$ and $\hat{\text{var}}(\hat{\beta}_j)$ is computed from (1.5). Note that as m approaches infinity (or $m \uparrow \infty$), the confidence interval lengths go to zero ($t_v \downarrow z$ where z denotes the standard normal variable, $\hat{\text{var}}(\hat{\beta}_j)(m-1) = \text{constant}$, $T_r \downarrow 0$). Rao (1959, p. 57) states that " C [in the example] is the least squares variance of the estimator"; actually we showed that C equals m times that variance! We can rewrite (3.1.b) as follows

$$\hat{\beta} \pm t_v \hat{\sigma}(\hat{\beta}_j) \left[\frac{(m-1)(1+T_r)}{(m-1)-(n-Q)} \right]^{\frac{1}{2}} \quad (3.1.c)$$

where $\hat{\sigma}(\hat{\beta}_j)$ denotes $\{\hat{\text{var}}(\hat{\beta}_j)\}^{\frac{1}{2}}$. In a saturate design we have $n = Q$; in practice n will not exceed Q very much. So assuming $(n-Q) \downarrow 0$ yields

$$\hat{\beta} \pm t_v \hat{\sigma}(\hat{\beta}_j) (1+T_r)^{\frac{1}{2}} \quad (3.2)$$

Eq. (2.4.b) proved that $T_r \downarrow 0$ if the regression model fits adequately or if there are many replications. Kleijnen (1988, p. 68) proposes to use EGLS only if $m \geq 25$ so that $\hat{\sigma}(\hat{\beta}_j)$ may be computed from (1.5); if $m \geq 25$ then t_v may be replaced by the standard normal variable z :

$$\hat{\beta} \pm z \hat{\sigma}(\hat{\beta}_j) . \quad (3.3)$$

4. A Monte Carlo experiment

We use a Monte Carlo experiment to estimate the α (or type I) and β (or type II) error of Rao's and Kleijnen's validation tests. We investigate Kleijnen's test only for the OLS estimator $\hat{\beta}$; see (1.2) and (1.4). [One reason for this restriction is that for the EGLS estimator $\hat{\beta}$ we know only the asymptotic covariance matrix; see (1.5). We expect that Kleijnen's test using EGLS will have a smaller β error (more power) than Kleijnen's test using OLS.] We further test the Studentized error using a t_v statistic with $v = m-1$ degrees of freedom. [Kleijnen (1983, p. 139) uses the standard normal statistic: $z = t_\infty$. Obviously $z^\alpha < t_v^\alpha$ for $v < \infty$ so that usage of z^α leads to higher type I errors and consequently to lower β errors.]

To estimate the actual α and β errors we apply Rao's and Kleijnen's tests 100 times per "case"; a case is defined by X , β , Q and m . The factor "100" means that an estimated $\hat{\alpha}$ is a binomial variable with standard error $\{\hat{\alpha}(1-\hat{\alpha})/100\}^{\frac{1}{2}}$; for example, if $\hat{\alpha} = 0.20$ (we take a nominal α of 0.20) then its standard error is 0.04. Likewise the estimated power $1 - \hat{\beta}$ has the standard error $\{\hat{\beta}(1-\hat{\beta})/100\}^{\frac{1}{2}}$; this error reaches its maximum value of 0.05 for $\hat{\beta} = 0.50$. To estimate β we must specify an alternative

model (or hypothesis H_1). We limit the Monte Carlo experiment to two factors x_2 and x_3 besides the dummy variable $x_1 = 1$; so we have

$$H_0 : E(\bar{y}_i) = \beta_1 + \beta_2 x_{i2} + \beta_3 x_{i3} \quad (4.1)$$

and we add a two-factor interaction β_4 to get

$$H_1 : E(\bar{y}_i) = \beta_1 + \beta_2 x_{i2} + \beta_3 x_{i3} + \beta_4 x_{i2} x_{i3} . \quad (4.2)$$

For X we select orthogonal and non-orthogonal matrices; see Table 3. (Note that the equations in the preceding sections all use X excluding the last column of X in Table 3, that is, $Q = 3$). Note that we select β_4 small relative to the "main effects" β_2 and β_3 . We fix $m = 10$. The response variances $\text{var}(y_i) = \sigma_i^2 = \sigma_{ii}$ are small for some cases (for example, case 1) and large for other cases (for example, case 2). For simplicity's sake we assume that the use of common random numbers in simulation yields constant correlation coefficients: $\rho_{ii'} = \sigma_{ii'}/(\sigma_i \sigma_{i'}) = \rho$ (a constant ρ was also assumed by several other authors; see Kleijnen, 1988). We investigate the effects of increasing correlation coefficients (more effective common seeds). If $\rho = 0$ then we might adapt Rao's approach; if we use different seeds in simulation then we know that $\rho = 0$ and we might replace the estimates $\hat{\sigma}_{ii'}$ in (1.1) by $\hat{\sigma}_{ii'} = 0$ for $i \neq i'$. This variant is shown in Tables 4 and 5 in the columns 0'. All results for case 1 are correlated since they use the same seed; they are independent of case 2, etc. (so rows 1 and 2 in Tables 4 and 5 are independent of all other rows).

TABLE 3
Monte Carlo Inputs

case	x_1 x_2 x_3 ($x_2 x_3$)	β_1 β_2 β_3 β_4	σ_1^2 ... σ_n^2
1)	$\begin{vmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{vmatrix}$	(1,10,10,1)	(1,1,1,1)
2)	same	(1,10,10,5)	(10,20,30,40)
3)	same	(1,10,20,1)	(1,1,1,1)
4)	same	(1,10,20,3)	(10,20,30,40)
5)	$\begin{vmatrix} 1 & -1 & 4 & -4 \\ 1 & 2 & -3 & -6 \\ 1 & -3 & 2 & -6 \\ 1 & 4 & -1 & -4 \\ 1 & 1 & 1 & 1 \end{vmatrix}$	(1,10,10,0.5)	(1,1,1,1,1)
6)	same	(1,10,10,3)	10.(1,2,3,4,5)
7)	same	(1,10,20,0.5)	(1,1,1,1,1)
8)	same	(1,10,20,5)	10.(1,2,3,4,5)
9)	$\begin{vmatrix} 1 & 1 & 6 & 6 \\ 1 & 2 & -5 & -10 \\ 1 & 3 & 4 & 12 \\ 1 & 4 & -3 & -12 \\ 1 & 5 & 2 & 10 \\ 1 & 6 & -1 & -6 \end{vmatrix}$	(1,10,10,0.25)	(1,1,1,1,1,1)
10)	same	(1,10,10,0.25)	10.(1,2,3,4,5,6)
11)	same	(1,10,20,0.25)	(1,1,1,1,1,1)
12)	same	(1,10,20,0.25)	10.(1,2,3,4,5,6)
13)	$\begin{vmatrix} 1 & 2 & -1 & -2 \\ 1 & 4 & -1 & -4 \\ 1 & 6 & 1 & 6 \\ 1 & 8 & 1 & 8 \\ 1 & 10 & -1 & -10 \\ 1 & 12 & -1 & -12 \end{vmatrix}$	(1,10,10,0.25)	(1,1,1,1,1,1)
14)	same	(1,10,10,0.5)	10.(1,2,3,4,5,6)
15)	same	(1,10,20,0.75)	(1,1,1,1,1,1)
16)	same	(1,10,20,1)	10.(1,2,3,4,5,6)

TABLE 4
The Estimated α Error: $\hat{\alpha}$ (nominal α is 0.20)

Case		0'	$\rho = 0$	0.1	0.3	0.5	0.7	0.9
1	Rao Kleijnen	0.20	0.22 0.01	0.22 0.01	0.22 0.01	0.25 0.00	0.24 0.00	0.20 0.00
2	Rao Kleijnen	0.10	0.14 0.03	0.13 0.03	0.15 0.01	0.13 0.00	0.11 0.00	0.12 0.00
3	Rao Kleijnen	0.15	0.16 0.03	0.16 0.01	0.17 0.00	0.18 0.00	0.18 0.00	0.19 0.00
4	Rao Kleijnen	0.17	0.21 0.02	0.21 0.01	0.21 0.01	0.20 0.00	0.19 0.00	0.19 0.00
5	Rao Kleijnen	0.14	0.23 0.11	0.22 0.09	0.21 0.04	0.21 0.01	0.21 0.00	0.20 0.00
6	Rao Kleijnen	0.06	0.13 0.03	0.12 0.02	0.15 0.04	0.16 0.03	0.15 0.00	0.18 0.00
7	Rao Kleijnen	0.17	0.17 0.11	0.17 0.08	0.23 0.04	0.24 0.00	0.23 0.00	0.21 0.00
8	Rao Kleijnen	0.11	0.14 0.08	0.13 0.05	0.15 0.04	0.16 0.00	0.16 0.00	0.17 0.00
9	Rao Kleijnen	0.09	0.19 0.13	0.20 0.12	0.19 0.06	0.17 0.04	0.17 0.01	0.15 0.00
10	Rao Kleijnen	0.08	0.19 0.09	0.17 0.09	0.16 0.08	0.16 0.04	0.16 0.02	0.18 0.00
11	Rao Kleijnen	0.05	0.19 0.08	0.18 0.07	0.19 0.05	0.18 0.04	0.19 0.00	0.18 0.00
12	Rao Kleijnen	0.10	0.18 0.09	0.18 0.09	0.18 0.06	0.16 0.03	0.16 0.00	0.14 0.00
13	Rao Kleijnen	0.11	0.26 0.16	0.27 0.13	0.28 0.07	0.29 0.03	0.28 0.00	0.24 0.00
14	Rao Kleijnen	0.06	0.19 0.15	0.17 0.11	0.17 0.07	0.17 0.04	0.16 0.00	0.16 0.00
15	Rao Kleijnen	0.14	0.20 0.17	0.22 0.17	0.25 0.09	0.26 0.05	0.25 0.00	0.27 0.00
16	Rao Kleijnen	0.08	0.22 0.10	0.21 0.08	0.21 0.06	0.20 0.02	0.21 0.00	0.25 0.00

Next we consider the results $\hat{\alpha}$ in Table 4. If Rao's derivations (including use of asymptotic results) are correct, then the following null-hypothesis (denoted by H_0^R to avoid confusion with H_0 in 4.1) holds:

$$H_0^R : E(\hat{\alpha}) = \alpha \quad (4.2.a)$$

and the alternative hypothesis is

$$H_1^R : E(\hat{\alpha}) \neq \alpha . \quad (4.2.b)$$

Kleijnen's approach uses the Bonferroni inequality so that we expect it to be conservative:

$$H_0^K : E(\hat{\alpha}) \leq \alpha \quad (4.3.a)$$

and

$$H_1^K : E(\hat{\alpha}) > \alpha . \quad (4.3.b)$$

We have 16 cases with independent seeds, each case yielding one binomial variable $\hat{\alpha}$ for a particular ρ . Under H_0^R and H_0^K respectively, we may add the 16 binomial variates to get a single binomial variate based on a sample size of 1600. The resulting binomial test per ρ value does not lead to rejection of H_0^R or H_0^K , except for our adaptation of Rao (column 0'); then $\hat{\alpha}$ is significantly low. In other words, Rao's method yields the correct α error; Kleijnen's method is conservative indeed.

Now we turn to the estimated power $1 - \hat{\beta}$ in Table 5. A casual inspection of the data suggests that Rao's power is higher. Indeed Wilcoxon's rank test for a bivariate sample $(\hat{\beta}^R, \hat{\beta}^K)$ shows that Rao's test has significantly higher power; our adaptation of Rao (column 0') has non-significantly lower power than Kleijnen's test.

TABLE 5
The Estimated Power $1 - \hat{\beta}$

Case		0'	$\rho = 0$	0.1	0.3	0.5	0.7	0.9
1	Rao Kleijnen	0.70	0.75 0.39	0.82 0.40	0.83 0.40	0.92 0.36	0.98 0.37	1.00 0.32
2	Rao Kleijnen	0.79	0.79 0.44	0.82 0.43	0.91 0.44	0.90 0.43	0.98 0.39	1.00 0.39
3	Rao Kleijnen	0.74	0.76 0.43	0.79 0.42	0.82 0.38	0.90 0.36	0.98 0.36	1.00 0.34
4	Rao Kleijnen	0.54	0.55 0.21	0.57 0.18	0.61 0.14	0.81 0.09	0.87 0.06	0.97 0.00
5	Rao Kleijnen	0.66	0.73 0.60	0.75 0.56	0.84 0.58	0.91 0.53	0.97 0.52	1.00 0.61
6	Rao Kleijnen	0.68	0.73 0.57	0.75 0.55	0.79 0.52	0.88 0.51	0.79 0.48	1.00 0.51
7	Rao Kleijnen	0.67	0.75 0.59	0.75 0.60	0.85 0.61	0.96 0.60	0.99 0.59	1.00 0.64
8	Rao Kleijnen	0.96	0.98 0.94	0.99 0.94	1.00 0.95	1.00 0.93	1.00 0.94	1.00 0.98
9	Rao Kleijnen	0.67	0.81 0.73	0.83 0.74	0.88 0.72	0.94 0.69	0.99 0.63	1.00 0.57
10	Rao Kleijnen	0.10	0.21 0.14	0.20 0.12	0.22 0.07	0.20 0.04	0.23 0.02	0.38 0.00
11	Rao Kleijnen	0.71	0.79 0.77	0.80 0.77	0.87 0.71	0.94 0.66	1.00 0.64	1.00 0.58
12	Rao Kleijnen	0.12	0.20 0.15	0.19 0.12	0.19 0.08	0.19 0.04	0.21 0.03	0.41 0.00
13	Rao Kleijnen	0.12	0.29 0.17	0.29 0.14	0.32 0.09	0.40 0.04	0.45 0.01	0.68 0.00
14	Rao Kleijnen	0.06	0.21 0.15	0.21 0.11	0.20 0.07	0.20 0.04	0.19 0.01	0.21 0.00
15	Rao Kleijnen	0.47	0.57 0.58	0.60 0.55	0.65 0.54	0.78 0.54	0.94 0.54	1.00 0.54
16	Rao Kleijnen	0.10	0.30 0.15	0.30 0.10	0.30 0.09	0.29 0.04	0.33 0.00	0.41 0.00

Summarizing so far, we see that Kleijnen's test devised for $\rho = 0$, is conservative ($E(\hat{\alpha}) < \alpha$) and (consequently) has lower power.

What is the effect of ρ ? Kleijnen's test was devised for $\rho = 0$ (independent seeds). Table 4 suggests that for $\rho \uparrow 1$ we have $E(\hat{\alpha}) \downarrow 0$. Indeed we may fit the following model to the six observations per case:

$$E(\hat{\alpha}_l) = \gamma_0 + \gamma_1 \rho_l \quad (l = 1, \dots, 6) \quad (4.4)$$

and $\hat{\gamma}_1$ is negative in each of the 16 cases. To understand this effect of ρ on $\hat{\alpha}$ we consider an extreme case: $\rho = 1$. Then all observations \bar{y}_i lie on a straight line ($\bar{y}_i = \beta_1 + \beta_2 x_{i2} + \beta_3 x_{i3}$); deleting one observation i does not affect $\hat{\beta}$; hence $\hat{\gamma}_i$ does not change (and remains equal to \bar{y}_i); and we do not reject the model. A smaller $\hat{\alpha}$ means lower power $1 - \hat{\beta}$; see Table 5. So we should not apply Kleijnen's method in case of common seeds.

What is the effect of ρ on Rao's test? We have already seen that Rao's type I error remains α (see Table 4). Table 5 suggests that increasing ρ increases the power. If $\rho = 0.9$ then the power is often 100%. Indeed fitting a model similar to (4.4) gives a positive slope ($\hat{\gamma}_1 > 0$) except for case 14.

5. Conclusions

If simulation experiments use common random numbers as inputs, then the simulation outputs y are correlated. To analyze the simulation data, OLS and EGLS can be applied; EGLS seems to require many replications; see Kleijnen (1988). When EGLS is applied, Rao (1959) provides a lack of fit test which is better than Kleijnen's cross-validation test. If the EGLS regression (meta)model is not rejected, then confidence intervals for individual effects β_j can be based on either Rao (1959) or Kleijnen (1988) which differ only slightly.

Acknowledgement

The Monte Carlo experiment of Section 4 was programmed by B. Annink and L. Simons, graduate students in the School of Business and Economics, Catholic University Brabant, Tilburg.

References

- Kleijnen, J.P.C., "Cross-Validation using the t Statistic", European J. Oper. Res., 13, 2(1983), 133-141.
- Kleijnen, J.P.C., Statistical Tools for Simulation Practitioners, Marcel Dekker, Inc., New York, 1987.
- Kleijnen, J.P.C., Analyzing Simulation Experiment with Common Random Numbers, Management Science, 34, 1(1988), 65-74.
- Rao, C.R., Some problems involving linear hypotheses in multivariate analysis, BIOMETRIKA, 46, (1959), 49-58.

IN 1987 REEDS VERSCHENEN

- 242 Gerard van den Berg
Nonstationarity in job search theory
- 243 Annie Cuyt, Brigitte Verdonk
Block-tridiagonal linear systems and branched continued fractions
- 244 J.C. de Vos, W. Vervaat
Local Times of Bernoulli Walk
- 245 Arie Kapteyn, Peter Kooreman, Rob Willemse
Some methodological issues in the implementation
of subjective poverty definitions
- 246 J.P.C. Kleijnen, J. Kriens, M.C.H.M. Lafleur, J.H.F. Pardoel
Sampling for Quality Inspection and Correction: AOQL Performance
Criteria
- 247 D.B.J. Schouten
Algemene theorie van de internationale conjuncturele en structurele
afhankelijkheden
- 248 F.C. Bussemaker, W.H. Haemers, J.J. Seidel, E. Spence
On (v,k,λ) graphs and designs with trivial automorphism group
- 249 Peter M. Kort
The Influence of a Stochastic Environment on the Firm's Optimal Dyna-
mic Investment Policy
- 250 R.H.J.M. Gradus
Preliminary version
The reaction of the firm on governmental policy: a game-theoretical
approach
- 251 J.G. de Gooijer, R.M.J. Heuts
Higher order moments of bilinear time series processes with symmetri-
cally distributed errors
- 252 P.H. Stevers, P.A.M. Versteijne
Evaluatie van marketing-activiteiten
- 253 H.P.A. Mulders, A.J. van Reeken
DATAAL - een hulpmiddel voor onderhoud van gegevensverzamelingen
- 254 P. Kooreman, A. Kapteyn
On the identifiability of household production functions with joint
products: A comment
- 255 B. van Riel
Was er een profit-squeeze in de Nederlandse industrie?
- 256 R.P. Gilles
Economies with coalitional structures and core-like equilibrium con-
cepts

- 257 P.H.M. Ruys, G. van der Laan
Computation of an industrial equilibrium
- 258 W.H. Haemers, A.E. Brouwer
Association schemes
- 259 G.J.M. van den Boom
Some modifications and applications of Rubinstein's perfect equilibrium model of bargaining
- 260 A.W.A. Boot, A.V. Thakor, G.F. Udell
Competition, Risk Neutrality and Loan Commitments
- 261 A.W.A. Boot, A.V. Thakor, G.F. Udell
Collateral and Borrower Risk
- 262 A. Kapteyn, I. Woittiez
Preference Interdependence and Habit Formation in Family Labor Supply
- 263 B. Bettonvil
A formal description of discrete event dynamic systems including perturbation analysis
- 264 Sylvester C.W. Eijffinger
A monthly model for the monetary policy in the Netherlands
- 265 F. van der Ploeg, A.J. de Zeeuw
Conflict over arms accumulation in market and command economies
- 266 F. van der Ploeg, A.J. de Zeeuw
Perfect equilibrium in a model of competitive arms accumulation
- 267 Aart de Zeeuw
Inflation and reputation: comment
- 268 A.J. de Zeeuw, F. van der Ploeg
Difference games and policy evaluation: a conceptual framework
- 269 Frederick van der Ploeg
Rationing in open economy and dynamic macroeconomics: a survey
- 270 G. van der Laan and A.J.J. Talman
Computing economic equilibria by variable dimension algorithms: state of the art
- 271 C.A.J.M. Dirven and A.J.J. Talman
A simplicial algorithm for finding equilibria in economies with linear production technologies
- 272 Th.E. Nijman and F.C. Palm
Consistent estimation of regression models with incompletely observed exogenous variables
- 273 Th.E. Nijman and F.C. Palm
Predictive accuracy gain from disaggregate sampling in arima - models

- 274 Raymond H.J.M. Gradus
The net present value of governmental policy: a possible way to find the Stackelberg solutions
- 275 Jack P.C. Kleijnen
A DSS for production planning: a case study including simulation and optimization
- 276 A.M.H. Gerards
A short proof of Tutte's characterization of totally unimodular matrices
- 277 Th. van de Klundert and F. van der Ploeg
Wage rigidity and capital mobility in an optimizing model of a small open economy
- 278 Peter M. Kort
The net present value in dynamic models of the firm
- 279 Th. van de Klundert
A Macroeconomic Two-Country Model with Price-Discriminating Monopolists
- 280 Arnoud Boot and Anjan V. Thakor
Dynamic equilibrium in a competitive credit market: intertemporal contracting as insurance against rationing
- 281 Arnoud Boot and Anjan V. Thakor
Appendix: "Dynamic equilibrium in a competitive credit market: intertemporal contracting as insurance against rationing"
- 282 Arnoud Boot, Anjan V. Thakor and Gregory F. Udell
Credible commitments, contract enforcement problems and banks: intermediation as credibility assurance
- 283 Eduard Ponds
Wage bargaining and business cycles a Goodwin-Nash model
- 284 Prof.Dr. hab. Stefan Mynarski
The mechanism of restoring equilibrium and stability in polish market
- 285 P. Meulendijks
An exercise in welfare economics (II)
- 286 S. Jørgensen, P.M. Kort, G.J.C.Th. van Schijndel
Optimal investment, financing and dividends: a Stackelberg differential game
- 287 E. Nijssen, W. Reijnders
Privatisering en commercialisering; een oriëntatie ten aanzien van verzelfstandiging
- 288 C.B. Mulder
Inefficiency of automatically linking unemployment benefits to private sector wage rates

- 289 M.H.C. Paardekooper
A Quadratically convergent parallel Jacobi process for almost diagonal matrices with distinct eigenvalues
- 290 Pieter H.M. Ruys
Industries with private and public enterprises
- 291 J.J.A. Moors & J.C. van Houwelingen
Estimation of linear models with inequality restrictions
- 292 Arthur van Soest, Peter Kooreman
Vakantiebestemming en -bestedingen
- 293 Rob Alessie, Raymond Gradus, Bertrand Melenberg
The problem of not observing small expenditures in a consumer expenditure survey
- 294 F. Boekema, L. Oerlemans, A.J. Hendriks
Kansrijkheid en economische potentie: Top-down en bottom-up analyses
- 295 Rob Alessie, Bertrand Melenberg, Guglielmo Weber
Consumption, Leisure and Earnings-Related Liquidity Constraints: A Note
- 296 Arthur van Soest, Peter Kooreman
Estimation of the indirect translog demand system with binding non-negativity constraints

IN 1988 REEDS VERSCHENEN

- 297 Bert Bettonvil
Factor screening by sequential bifurcation
- 298 Robert P. Gilles
On perfect competition in an economy with a coalitional structure
- 299 Willem Selen, Ruud M. Heuts
Capacitated Lot-Size Production Planning in Process Industry
- 300 J. Kriens, J.Th. van Lieshout
Notes on the Markowitz portfolio selection method
- 301 Bert Bettonvil, Jack P.C. Kleijnen
Measurement scales and resolution IV designs: a note
- 302 Theo Nijman, Marno Verbeek
Estimation of time dependent parameters in linear models
using cross sections, panels or both
- 303 Raymond H.J.M. Gradus
A differential game between government and firms: a non-cooperative
approach
- 304 Leo W.G. Strijbosch, Ronald J.M.M. Does
Comparison of bias-reducing methods for estimating the parameter in
dilution series
- 305 Drs. W.J. Reijnders, Drs. W.F. Verstappen
Strategische bespiegelingen betreffende het Nederlandse kwaliteits-
concept
- 306 J.P.C. Kleijnen, J. Kriens, H. Timmermans and H. Van den Wildenberg
Regression sampling in statistical auditing
- 307 Isolde Woittiez, Arie Kapteyn
A Model of Job Choice, Labour Supply and Wages
- 308 Jack P.C. Kleijnen
Simulation and optimization in production planning: A case study
- 309 Robert P. Gilles and Pieter H.M. Ruys
Relational constraints in coalition formation
- 310 Drs. H. Leo Theuns
Determinanten van de vraag naar vakantiereizen: een verkenning van
materiële en immateriële factoren
- 311 Peter M. Kort
Dynamic Firm Behaviour within an Uncertain Environment
- 312 J.P.C. Blanc
A numerical approach to cyclic-service queueing models

- 313 Drs. N.J. de Beer, Drs. A.M. van Nunen, Drs. M.O. Nijkamp
Does Morkmon Matter?
- 314 Th. van de Klundert
Wage differentials and employment in a two-sector model with a dual labour market
- 315 Aart de Zeeuw, Fons Groot, Cees Withagen
On Credible Optimal Tax Rate Policies
- 316 Christian B. Mulder
Wage moderating effects of corporatism
Decentralized versus centralized wage setting in a union, firm, government context
- 317 Jörg Glombowski, Michael Krüger
A short-period Goodwin growth cycle
- 318 Theo Nijman, Marno Verbeek, Arthur van Soest
The optimal design of rotating panels in a simple analysis of variance model
- 319 Drs. S.V. Hannema, Drs. P.A.M. Versteijne
De toepassing en toekomst van public private partnership's bij de grote en middelgrote Nederlandse gemeenten
- 320 Th. van de Klundert
Wage Rigidity, Capital Accumulation and Unemployment in a Small Open Economy
- 321 M.H.C. Paardekooper
An upper and a lower bound for the distance of a manifold to a nearby point
- 322 Th. ten Raa, F. van der Ploeg
A statistical approach to the problem of negatives in input-output analysis
- 323 P. Kooreman
Household Labor Force Participation as a Cooperative Game; an Empirical Model
- 324 A.B.T.M. van Schaik
Persistent Unemployment and Long Run Growth
- 325 Dr. F.W.M. Boekema, Drs. L.A.G. Oerlemans
De lokale produktiestructuur doorgelicht.
Bedrijfstakverkenningen ten behoeve van regionaal-economisch onderzoek
- 326 J.P.C. Kleijnen, J. Kriens, M.C.H.M. Lafleur, J.H.F. Pardoel
Sampling for quality inspection and correction: AOQL performance criteria

- 327 Theo E. Nijman, Mark F.J. Steel
Exclusion restrictions in instrumental variables equations
- 328 B.B. van der Genugten
Estimation in linear regression under the presence of heteroskedasticity of a completely unknown form
- 329 Raymond H.J.M. Gradus
The employment policy of government: to create jobs or to let them create?
- 330 Hans Kremers, Dolf Talman
Solving the nonlinear complementarity problem with lower and upper bounds
- 331 Antoon van den Elzen
Interpretation and generalization of the Lemke-Howson algorithm

Bibliotheek K. U. Brabant



17 000 01065919 2